

Aberystwyth University

Multiple melt bodies fed the AD 2011 eruption of Puyehue-Cordon Caulle, Chile

Alloway, B. V.; Pearce, N. J. G.; Villarosa, G.; Outes, V.; Moreno, P. I.

Published in:
Scientific Reports

DOI:
[10.1038/srep17589](https://doi.org/10.1038/srep17589)

Publication date:
2015

Citation for published version (APA):

Alloway, B. V., Pearce, N. J. G., Villarosa, G., Outes, V., & Moreno, P. I. (2015). Multiple melt bodies fed the AD 2011 eruption of Puyehue-Cordon Caulle, Chile. *Scientific Reports*, 5, [17589].
<https://doi.org/10.1038/srep17589>

Document License CC BY

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

SCIENTIFIC REPORTS

OPEN

Multiple melt bodies fed the AD 2011 eruption of Puyehue-Cordón Caulle, Chile

B. V. Alloway¹, N. J. G. Pearce², G. Villarosa³, V. Outes³ & P. I. Moreno⁴

Received: 27 May 2015

Accepted: 02 November 2015

Published: 02 December 2015

Within the volcanological community there is a growing awareness that many large- to small-scale, point-source eruptive events can be fed by multiple melt bodies rather than from a single magma reservoir. In this study, glass shard major- and trace-element compositions were determined from tephra systematically sampled from the outset of the Puyehue-Cordón Caulle (PCC) eruption (~1 km³) in southern Chile which commenced on June 4th, 2011. Three distinct but cogenetic magma bodies were simultaneously tapped during the paroxysmal phase of this eruption. These are readily identified by clear compositional gaps in CaO, and by Sr/Zr and Sr/Y ratios, resulting from dominantly plagioclase extraction at slightly different pressures, with incompatible elements controlled by zircon crystallisation. Our results clearly demonstrate the utility of glass shard major- and trace-element data in defining the contribution of multiple magma bodies to an explosive eruption. The complex spatial association of the PCC fissure zone with the Liquiñe-Ofqui Fault zone was likely an influential factor that impeded the ascent of the parent magma and allowed the formation of discrete melt bodies within the sub-volcanic system that continued to independently fractionate.

At 18.45 UTC on June 4th, 2011, an explosive eruption (VEI ≤ 5) of rhyolitic magma^{1–3} began at Puyehue-Cordón Caulle volcanic complex (PCCVC), southern Chile and close to the frontier with Argentina. This PCC eruption was the first within this dominantly dacitic to rhyolitic fissure zone since 1960^{4–7} (Fig. 1) and was preceded (commencing April 27th, 2011) by intensifying seismic swarms located between 2.5 and 5 km depth¹. The eruption commenced with the development of a sustained Plinian eruption column that ascended to >15 km above the newly formed vent. Ash from this column circum-navigated the southern Hemisphere in less than 14 days and severely disrupted aviation in Argentina, Uruguay, South Africa, Australia, New Zealand and finally in Chile itself. By January 2012, explosive activity had declined from sustained Plinian to intermittent sub-Plinian fountaining, to mixed gas- and ash-jetting punctuated by Vulcanian blasts. This explosive activity was accompanied by synchronous effusion of obsidian lava in a hybrid explosive-effusive eruption^{1,3}. Throughout the PCC eruption two closely spaced (<200 m), and sometimes partly overlapping, explosive loci (NW and SE vents) were noted³. The regular occurrence of synchronous blasts from both vents suggested an advanced degree of connectivity in the shallow subsurface.

The position of the 2011 PCC vent(s) coincides with the intersection of the N-S oriented Liquiñe-Ofqui Fault zone with the 15-km long by 4 km wide NW-SE oriented Cordón Caulle fault-bounded graben with associated fissure vents, aligned domes and pyroclastic cones (Fig. 1).

In this study, we determine the major- and trace-element chemistries of fresh vesicular glass shards using grain-discrete electron microprobe (EMP) and laser ablation inductively coupled plasma-mass spectrometry (LA-ICP-MS) techniques, respectively. Daily ash-fall samples were collected from downwind

¹School of Geography, Environment and Earth Sciences, Victoria University of Wellington, Wellington 6140, New Zealand. ²Department of Geography & Earth Sciences, Aberystwyth University, SY23 3DB Wales, United Kingdom.

³INIBIOMA, CONICET-Universidad Nacional del Comahue, Bariloche, Argentina. ⁴Department of Ecological Sciences and Millennium Institute of Ecology and Biodiversity, Universidad de Chile, Santiago 7750000, Chile. Correspondence and requests for materials should be addressed to B.V.A. (email: brent.alloway@vuw.ac.nz)




Figure 1. Map of the southeast portion of the PCC volcanic complex showing the location and spatial arrangement of 1921 (coloured orange), 1960 (coloured red) and 2011 (coloured yellow) fissure vents and resultant lava products (inset modified from *ref. 5*). Surveyed points are indicated in meters above sea level. Maps were modified using Adobe Illustrator CS5 version 15.0.2.

localities (*SI Table 1*) from the commencement of the PCC eruption on June 4th, 2011 in order to quantify any compositional evolution of the rhyolitic magma as the eruption progressed. Fresh vesicular glass was specifically selected during analysis to minimize potential contamination by conduit and/or overburden glassy materials that may have been incorporated during magma ascent.

Methods

PCC ash collection and isopach map. Ash was systematically measured and collected from planar surfaces and collection receptacles from the outset of the PCC eruption on June 4th following a set of instructions^{8,9} adapted from IVHHN and USGS procedures. Based on this extensive collection network of 400 sampling points, which spanned the western half of the Argentine Provinces of Neuquén and Rio Negro, we produced a modified isopach map that indicates total composite ash thickness deposited during the paroxysmal phase of the eruption (Fig. 2A) with a volume of $\sim 1 \text{ km}^3$ (see also *ref. 10*). We also utilized NASA MODIS satellite imagery taken during the PCC eruption to monitor the plume during June 2011. Our isopach data highlights the complexity of mapping this variably directed and widely dispersed PCC plume by a sustained eruption of relatively small volume (Fig. 2B). On this basis, it is clear that this PCC eruption shares many characteristics with that of the equally small-volume A.D. 2010 Icelandic eruption of Eyjafjallajökull Volcano that created significant disruption across mainland Europe due to its rapid dispersal by prevailing westerly winds^{11,12}.

Electron Microprobe (EMP) technique. Major element determinations were made on a JEOL Superprobe (JXA-8230) housed at Victoria University of Wellington, using the ZAF correction method. Analyses were performed with 15kV accelerating voltage, 8 nA beam current, and an electron beam



defocused to between 20 to 10 μm . Standardization was achieved by means of mineral and glass standards. A rhyolitic glass standard (VG-568) was routinely used to monitor calibration in all analytical runs, and used to evaluate any day-to-day differences in the calibration. The large number of samples precluded conducting all analyses in a single batch. All analyses are normalized to 100 wt.% anhydrous, with H_2O by difference being given, and total Fe is reported as FeO. Glass shard major element analyses are presented in *SI Table 2*.

Laser ablation inductively coupled plasma-mass spectrometry (LA-ICP-MS) technique. Trace element analyses on individual glass shards were performed by laser ablation (LA) ICP-MS in the Department of Geography and Earth Sciences, Aberystwyth University, using a Coherent GeoLas ArF 193 nm Excimer LA system coupled to a Thermo Finnegan Element 2 sector field ICP-MS. Trace element data were collected for individual shards with the majority of analyses performed using 20 μm ablation craters. Laser fluence was 10 Jcm⁻² at a repetition rate of 5 Hz for a 24 second acquisition. The minor ²⁹Si isotope was used as the internal standard, with SiO₂ (determined by EMPA) used to calibrate each analysis, after normalization to an anhydrous basis. The NIST 612 reference glass was used for calibration, taking concentrations from *ref.* 13. A fractionation factor was applied to the data to account for analytical bias related to the different matrices of the reference standard and the sample. For this factor

as well as ICP-MS and laser operating conditions see *ref.* 14, and references therein. The MPI-DING reference glass ATHO-G¹⁵ was analysed as an unknown under the same operating conditions at the same time. Analytical precision is typically between ± 5 –10%, and accuracy is typically around ± 5 %, when compared with the published GeoReM concentrations for ATHO-G. Glass shard trace element analyses are presented in *SI Table 3*.

X-ray fluorescence (XRF) and solution inductively coupled plasma mass spectrometry (ICP-MS) techniques. Bulk glass major and trace element compositions of PCC eruptive material were obtained by X-ray fluorescence (XRF) and solution inductively coupled plasma mass spectrometry (ICP-MS) analyses conducted at CRL Energy Ltd. SpectraChem Analytical facilities located in Lower Hutt, New Zealand, and at Victoria University of Wellington, New Zealand, respectively. Washed pumiceous lapilli were crushed and then sieved between 250- to 63- μm . Crystal and glass components were then separated using a Frantz magnetic separator, with the purity of the separated glass visually assessed using a binocular microscope. A total of 15 analyses were carried out: one sample (PCC-7) was run in duplicate and the BCR-2 basaltic rock standard was analysed to check accuracy. Bulk glass major (XRF) and trace element (Solution-ICP-MS) were determined using Siemens SRS3000 and Agilent 7500CS ICP-MS instruments, respectively, with samples bracketed by the primary standard (BCR-2). Analyses are presented in *SI Table 4*.

Geochemistry and Grain-Size Attributes

Major element analyses conducted on glass shards reveal three compositional groupings (Fig. 3A–C; *SI Table 2*) defined by minima in CaO concentrations at 1.57 and 1.37 wt% (Fig. 3D). These groups are present in each of the samples collected on different days indicating that they are not analytical artefacts. With increasing SiO_2 , there is a steady decrease in CaO, FeO, MgO, TiO_2 , Al_2O_3 and Na_2O , with a corresponding slight increase in K_2O , consistent with fractionation of sodic plagioclase, pyroxene and magnetite/ilmenite (cf. *ref.* 2). Projection of average group compositions (divided on CaO, see above) into the synthetic system $\text{Qz}'\text{-Ab}'\text{-Or}'$ (following *ref.* 16) shows evolution of the magmas towards granite minimum compositions (Fig. 3E), but never reaching a cotectic or minimum composition. The magmatic evolution between the three groups instead suggests ascent-driven crystallisation with three separate pockets of magma becoming arrested and evolving at different depths in the sub-volcanic system and most likely, at pressures of between 100 to 50 MPa (see *ref.* 2, cf. *ref.* 16). Indeed, these experimentally determined crystallisation pressures are in broad agreement with the 2.5- to 5-km depths of pre-eruption seismic swarms that commenced on April 27th, 2011.

Close inspection of the data shows a gradual decline of CaO with increasing SiO_2 within each group, superimposed on the general between-group trend of decreasing CaO with increasing SiO_2 . Trace element data (Fig. 4; *SI Table 3*) mirrors these major element trends, with Sr echoing the groupings based on CaO compositions, and Sr/Zr ratios clearly defining the three separate evolving magma batches (with Sr/Zr < 0.34, 0.34–0.39, and > 0.39). Variations in Zr, Y and Th result from minor zircon extraction within these groups concurrent with dominant plagioclase (and pyroxene) fractionation (Fig. 4A–C). Incompatible elements (including Nb, Y, Zr, REE, Hf, Ta, Th, U) show consistent element/element ratios attesting to the co-magmatic origin of the three magma groups (Fig. 4D–F). Once again, the trace element data (e.g. Sr vs. Zr, Y and Th) shows that each separate magmatic group is represented in each of the individual days' eruptive products (see Fig. 4), indicating the simultaneous tapping of these compositionally distinct melt bodies.

The bulk chemistry shows a trend of plagioclase extraction that defines a fractionation trend between a phenocryst assemblage dominated by plagioclase and the EPMA glass compositions (72.37 ± 0.60 wt.% SiO_2 ; 1.43 ± 0.08 wt.% CaO) (*SI Fig. 1*; *SI Table 4*). Incompatible trace element ratios are consistent between bulk sample and individual glass shards (particularly when the compatibility of some trace elements in phenocrysts is considered, e.g. Y in pyroxene), and these compare well with similarly acquired data (*SI Fig. 2*) from Puyehue Volcano and its adjacent volcanic massif⁷.

Laser auto-analyser grain-size analyses were also conducted on each tephra sample and indicate multimodal grain-size peaks (*SI Fig. 3*), though without conducting grain discrete geochemistry on glass shards from each of these peaks it is difficult to accurately assess the relationship of a particular grain-size component to a potential magmatic source. Irrespective of this, the key control in determining the grain-size characteristics appears to be dominantly influenced by the variable meteorological conditions affecting the ash column at different altitudes and on different days in combination with variations in eruptive flux¹⁰.

Discussion

Despite a number of studies^{9,10,17,18} that closely examined the products of the AD 2011 PCC eruption and, indeed, verified its eruptive complexity, the contribution of magma reservoirs and their subtle geochemical differences were not recognised. Our results clearly demonstrate the utility of grain discrete glass shard major- and trace-element data as a relatively simple and straightforward method by which the involvement of multiple magma bodies contributing to a point-source eruption can be easily detected. We suggest that the formation of separate pockets of intermediate to evolved magma, which arrest at different depths during ascent, and then become variously involved in a point-source eruptive event, is




Figure 3. (A–C) Selected major element compositions (weight percent SiO₂ vs. CaO and FeO vs. K₂O and CaO) of glass shards from PCC tephra collected in time series; (D) Frequency histogram of glass shard wt% CaO identifying compositional gaps of 1.37 and 1.57 wt% respectively; All PCC-glasses are peraluminous (molar Al₂O₃ > (Na₂O + K₂O + CaO)) and thus corundum normative; (E) Projection of the PCC glass compositional groups into the synthetic system Qz'-Ab'-Or' (see ref. 16). Selected H₂O saturated phase boundaries in Qz'-Ab'-Or' haplogranitic melts as a function of pressure are indicated.

likely to be a more common eruptive phenomenon than previously recognised (i.e. see ref. 19), especially in highly complex volcano-tectonic environments such as the Andes.

The involvement of multiple magma bodies during the paroxysmal phase of the AD 2011 PCC eruption may also partially explain why hybrid explosive-effusive activities were near-simultaneously produced³. Our results are consistent with earlier studies (i.e. ref. 6) suggesting that the highly fractured nature of the PCC fissure zone was influential in defining the tempo and style of volcanism within this zone. The complex setting of the PCCVC, adjacent to regional tectonic structures such as the Liquiñe-Ofqui Fault zone, coupled with its extant (pre-2011) magmatic plumbing system, is likely to be influential in allowing




Figure 4. Selected trace element plots of PCC samples. Sr vs. Zr, Y and Th (A–C) indicate three closely related compositional fields that correspond with major element compositional clusters identified in Fig. 3 (A–C). The PCC glasses are all peraluminous (corundum normative) rhyolites, containing between 300–400 ppm Zr, and thus will be zircon saturated²³. The variation of Sr in bulk samples (see insets A'–C') indicates that plagioclase fractionation is important in the evolution of the PCC magma, is consistent with both phenocryst observations and experiment (cf. ref. 2), and mirrors the major element data. In the glasses, Zr, Y and Th (A–C) (all highly compatible in zircon) decrease with decreasing Sr indicating the concurrent fractionation of zircon alongside plagioclase. Incompatible elements such as Y vs. Nb, Ce and Th (D–F) show no variation in element/element ratios between the three magma groups, indicating the cogenetic relationship of the simultaneously erupted magma bodies.

or impeding magma ascent, and thus potentially facilitating the formation of separate (independent, fractionating) melt bodies within the sub-volcanic system.

Recent petrological and InSAR measurements²⁰ propose a pre-existing laccolith-shaped magma reservoir at 4 to 7 km depth, with an inferred area of 20 km², which may have fed all three historical (1921, 1960 and 2011) eruptions. This extensive reservoir presumably 'leaked' magma upwards forming the three compositionally distinct melt bodies involved in the 2011 PCC eruption. How these melt bodies might have been temporally and spatially linked with each other and with this larger reservoir system and then simultaneously activated remains to be determined. The application of intra-crystalline diffusion

studies^{21,22} should eventually reveal more specific time-series changes in magmatic conditions prior to, during and after magma emplacement while analysis of pre- and syn-eruption InSAR and seismic data along with future geophysical transects across this fissure zone may locate and configure magma storage levels, assess interconnectivity as well as identify spatial relationships between surface and sub-surface fault and pre-2011 magmatic structures. Ultimately, these approaches will be useful in clarifying some of the unresolved questions posed from this research.

References

1. Silva Parejas, C. *et al.* The 2011–2012 eruption of Cordón Caulle volcano (southern Andes): evolution, crisis management and current hazards. *EGU general Assembly Abstracts* **14** (EGU2012-9382-2) (2012).
2. Castro, J. M. *et al.* Storage and eruption of near-liquidus rhyodacitic magma at Cordón Caulle. *Chile. Bull. Volcan.* **75**, 702 (2013).
3. Schipper, C. I. *et al.* Shallow vent architecture during hybrid explosive-effusive activity at Cordón Caulle (Chile, 2011–2012): Evidence from direct observations and pyroclast textures. *Journ. Volcan. Geotherm. Res.* **262**, 25–37 (2013).
4. Lara, L. E. & Moreno, H. Geología del Complejo Volcánico Puyehue-Cordón Caulle, X Región de Los Lagos. Servicio Nacional de Geología y Minería. Carta Geológica de Chile. Serie Geología Básica, 1 mapa escala 1:50,000 (2006).
5. Lara, L. E. *et al.* Rhyodacite fissure eruption in Southern Andes (Cordón Caulle; 40.5°S) after the 1960 (Mw: 9.5 Chilean earthquake: a structural interpretation. *Journ. Volcan. Geotherm. Res.* **138**, 127–138 (2004).
6. Lara, L. E. *et al.* Magmatic evolution of the Puyehue-Cordón Caulle Volcanic Complex (40°S), Southern Andean Volcanic Zone: from shield to unusual rhyolitic fissure volcanism. *Journ. Volcan. Geotherm. Res.* **157**, 343–366 (2006).
7. Singer, B. S. *et al.* Eruptive history, geochronology and magmatic evolution of the Puyehue-Cordon Caulle volcanic complex, Chile. *Geol. Soc. Amer. Bull.* **120**, 599–618 (2008).
8. Villarosa, G. & Outes, V. Rutina de toma de muestras de tefra: procedimientos para coleccionar muestras de ceniza. *INIBIOMA (CONICET-Universidad Nacional del Comahue) booklet*. **6** pp. (2008).
9. Collini, E. *et al.* Volcanic ash forecast during the June 2011 Cordón Caulle eruption. *Nat. Hazards* **66**, 389–412 (2013).
10. Pistolesi, M. *et al.* Complex dynamics of small-moderate volcanic events: the example of the 2011 rhyolitic Cordón Caulle eruption, Chile. *Bull. Volc.* **77**, no.3, doi: 10.1007/s00445-014-0898-3 (2015).
11. UK Met Office. Volcanic Ash Advisory Centre, London, UK. PFXD01_EGRR Advisory files from 1146 hrs, April 14th, 2010 (Advisory NR 2010/001) to 1146 hrs, May 24th, 2010 (Advisory NR 2010/156) http://www.metoffice.gov.uk/aviation/vaac/vaacuk_vag.html (2010). Date of access: 12/12/2014.
12. Davies, S. M. *et al.* Widespread dispersal of Icelandic tephra: How does the Eyjafjöll eruption of 2010 compare to past Icelandic events? *Journ. Quat. Sci.* **25**, 605–611 (2010).
13. Pearce, N. J. G. *et al.* A compilation of new and published major and trace element data for NIST SRM 610 and NIST SRM 612 glass reference materials. *Geost. Newslet.* **21**, 115–144 (1997).
14. Pearce, N. J. G. *et al.* Trace-element analysis by LA- ICP-MS: the quest for comprehensive chemical characterisation of single sub-10µm volcanic glass shards. *Quat. Int.* **246**, 57–81 (2011).
15. Jochum, K. P., Stoll, B. & Herwig, K. *et al.* MPI-DING reference glasses for *in situ* microanalysis: new reference values for element concentrations and isotope ratios. *Geochem. Geophys. Geosys.* **7**, Q02008, doi: 02010.01029/02005GC001060 (2006).
16. Blundy, J. & Cashman, K. Ascent driven crystallisation of dacite magmas at Mt. St. Helens, 1980–1986. *Contrib. Mineral. Petrol.* **140**, 631–650 (2001).
17. Bertrand, S. *et al.* Deposition of the 2011–2012 Cordón Caulle tephra (Chile, 40°S) in lake sediments: Implications for tephrochronology and volcanology. *Journ. Geophys. Res.: Earth Surface* **119**, 2555–2573, doi: 10.1002/2014JF003321 (2014).
18. Daga, R. *et al.* Characterisation of tephra dispersed by the recent eruptions of Calbuco (1961), Chaiten (2008) and Cordón Caulle Complex (1960 and 2011), in Northern Patagonia. *Journ. Sth Amer. Earth Sci.* **49**, 1–14, doi: 10.1016/j.jsames.2013.10.006 (2014).
19. Cashman, K. V. & Giordano, G. Calderas and magma reservoirs. *Journ. Volc. Geotherm. Res.* **288**, 28–45 (2014).
20. Jay, J. *et al.* Locating magma reservoirs using InSAR and petrology before and during the 2011–2012 Cordón Caulle silicic eruption. *Earth Planet. Sci. Lett.* **395**, 254–266 (2014).
21. Gualda G. A. R. *et al.* Timescales of quartz crystallization and the longevity of the Bishop giant magma body. *PLoS ONE* **7**, e37492, doi: 10.1371/journal.pone.0037492 (2012).
22. Saunders, K. *et al.* Linking Petrology and Seismology at an Active Volcano. *Science* **336**, 1023–1027, 10.1126/science.1220066 (2012).
23. Watson, E. B. Zircon saturation in felsic liquids: Experimental results and applications to trace element geochemistry. *Contrib. Mineral. Petrol.* **70**, 407–419 (1979).

Acknowledgements

This study was funded by a Victoria University of Wellington Science Faculty Research Grant (to BVA), Aberystwyth University Research Fund (to NJGP), Argentine PIP CONICET 2011 0311 and FONCYT PICT 2010 2046 (to GV), and Iniciativa Científica Milenio grants P02-51 and NC120066, Chilean Fondecyt 1151469 (to PIM). S.J. Cronin is thanked for organizing PCC ash leachate experiments at Massey University, New Zealand. We also thank Ms. Maria Noel Serra, CONICET, for ash collection field assistance and Ms. Ashley Pocock for technical assistance in the production of PCC tephra grain-size analyses conducted at Victoria University of Wellington.

Author Contributions

B.V.A. conducted the EMP analyses. B.V.A. and N.J.G.P. jointly conducted the LA-ICP-MS analyses and subsequent data analysis. G.V. and V.O. organized the collection of PCC tephra samples and produced the isopach map. PM provided support for field and laboratory activities. All authors were involved in the production of this paper.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Alloway, B. V. *et al.* Multiple melt bodies fed the AD 2011 eruption of Puyehue-Cordón Caulle, Chile. *Sci. Rep.* **5**, 17589; doi: 10.1038/srep17589 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>